arXiv:math/0305339v1 [math.NT] 23 May 2003

On the second moment of $S(T)$ in the theory of the Riemann zeta function

Tsz Ho Chan

February 1, 2008

Abstract

We assume the Riemann Hypothesis and an quantitative form of the Twin Prime Conjecture, and obtain an asymptotic formula for the second moment of $S(T)$ with better error term.

1 Introduction

Let $\rho = \beta + i\gamma$ be the zeros of the Riemann zeta function $\zeta(s)$. For $T \neq \gamma$,

$$S(T) = \frac{1}{\pi}\arg\zeta(\frac{1}{2} + iT),$$

where the argument is obtained by continuous variation along the horizontal line $\sigma + iT$ starting with the value zero at $\infty + iT$. For $T = \gamma$, we define

$$S(T) = \lim_{\epsilon \to 0} \frac{1}{2}\{S(T+\epsilon) + S(T-\epsilon)\}.$$

In [3], Goldston proved that under the Riemann Hypothesis,

$$\begin{aligned}\int_0^T |S(t)|^2 dt \quad = \quad & \frac{T}{2\pi^2}\log\log T + \frac{T}{2\pi^2}\Big[\int_1^\infty \frac{F(\alpha,T)}{\alpha^2} + C_0 \\ & + \sum_{m=2}^{\infty}\sum_p \Big(\frac{-1}{m} + \frac{1}{m^2}\Big)\frac{1}{p^m}\Big] + o(T)\,. \end{aligned} \tag{1}$$

Here C_0 is Euler's constant, and

$$F(\alpha) = F(\alpha,T) = \Big(\frac{T}{2\pi}\log T\Big)^{-1} \sum_{0<\gamma,\gamma'\leq T} T^{i\alpha(\gamma-\gamma')} w(\gamma-\gamma') \tag{2}$$

with $w(u) = 4/(4+u^2)$ is Montgomery's pair correlation function. Here and throughout this paper, p will denote a prime and sums over p are over all primes.

In [1] and [2], the author assumed the Riemann Hypthesis and the following quantitative form of Twin Prime Conjecture: For any $\epsilon > 0$,

$$\sum_{n=1}^{N} \Lambda(n)\Lambda(n+d) = \mathfrak{S}(d)N + O(N^{1/2+\epsilon})$$

uniformly in $|d| \leq N$. $\Lambda(n)$ is the von Mangoldt lambda function. $\mathfrak{S}(d) = 2\prod_{p>2}\big(1-1/(p-1)^2\big)\prod_{p|d,p>2}(p-1)/(p-2)$ if d is even, and $\mathfrak{S}(d) = 0$ if d is odd.

He proved that, for any $\epsilon > 0$,

$$F(\alpha, T) = \begin{cases} \alpha + \frac{1}{T^{2\alpha}}[\log T - 2\log 2\pi - 2] & \\ +O(\alpha T^{\alpha-1}) + O\Big(\frac{1}{T^{(1/2-\epsilon)\alpha}\log T}\Big), & \text{if } 0 \leq \alpha \leq 1 - \frac{3\log\log T}{\log T}, \\ \alpha + O(\frac{T^{\alpha-1}}{\log T}), & \text{if } 1 - \frac{3\log\log T}{\log T} \leq \alpha \leq 1. \end{cases} \tag{3}$$

Using this and more careful calculations, we have

Theorem 1.1. *Assume the Riemann Hypothesis and the above Twin Prime Conjecture. We have*

$$\begin{aligned} \int_0^T |S(t)|^2 dt &= \frac{T}{2\pi^2}\log\log T + \frac{T}{2\pi^2}\Big[\int_1^\infty \frac{F(\alpha,T)}{\alpha^2}d\alpha + C_0 \\ &\quad - \sum_{m=2}^{\infty}\sum_{p}\Big(\frac{1}{m} - \frac{1}{m^2}\Big)\frac{1}{p^m}\Big] + O\Big(\frac{T}{\log^2 T}\Big). \end{aligned}$$

So, there appears to be no $T/\log T$ term. However, one has to be careful about this because we still do not know anything very precise about

$$\int_1^\infty \frac{F(\alpha,T)}{\alpha^2}d\alpha.$$

Goldston [3] proved that under the Riemann Hypothesis, for any $\epsilon > 0$,

$$\frac{2}{3} - \epsilon < \int_1^\infty \frac{F(\alpha,T)}{\alpha^2}d\alpha < 2.$$

Montgomery [4] conjectured that

$$F(\alpha, T) = 1 + o(1) \text{ uniformly for } 1 \leq \alpha \leq M,$$

for any fixed M. This implies

$$\int_1^\infty \frac{F(\alpha,T)}{\alpha^2}d\alpha = 1 + o(1)$$

which gives an error term $o(T)$ for our Theorem 1.1. In order to get better error terms, one has to understand $F(\alpha, T)$ better for larger range of α, say $1 \leq \alpha \leq \log T$. This would be a great challenge.

2 Preparations

Essentially, our proof follows that of Goldston [3]. We shall recall some of his results.

Lemma 2.1. *Assume the Riemann Hypothesis. For* $t \geq 1$, $t \neq \gamma$, $x \geq 4$, *we have*

$$\begin{aligned} S(t) = &- \frac{1}{\pi} \sum_{n \leq x} \frac{\Lambda(n)}{n^{1/2}} \frac{\sin(t \log n)}{\log n} f\Big(\frac{\log n}{\log x}\Big) \\ &+ \frac{1}{\pi} \sum_{\gamma} \sin((t-\gamma)\log x) \int_0^\infty \frac{u}{u^2 + ((t-\gamma)\log x)^2} \frac{du}{\sinh u} \\ &+ O\Big(\frac{x^{1/2}}{t^2 \log x}\Big) + O\Big(\frac{1}{t \log x}\Big), \end{aligned} \tag{4}$$

where

$$f(u) = \frac{\pi}{2} u \cot\Big(\frac{\pi}{2} u\Big), \tag{5}$$

and $\Lambda(n) = \log p$ *if* $n = p^m$, *for* p *a prime and* $m \geq 1$, *and* $\Lambda(n) = 0$ *otherwise.*

Proof: This is Lemma 1 in [3].

Lemma 2.2. *For* $x \geq 4$ *and* $T \geq 2$,

$$\begin{aligned} R &= \int_1^T \Big| \frac{1}{\pi} \sum_{\gamma} \sin((t-\gamma)\log x) \int_0^\infty \frac{u}{u^2 + ((t-\gamma)\log x)^2} \frac{du}{\sinh u} \Big|^2 dt \\ &= \frac{1}{\pi^2 \log x} \sum_{\gamma, \gamma'} \hat{k}((\gamma - \gamma')\log x) + O(\log^3 T), \end{aligned}$$

where

$$k(u) = \begin{cases} \Big(\frac{1}{2u} - \frac{\pi^2}{2}\cot(\pi^2 u)\Big)^2, & \textit{if } |u| \leq \frac{1}{2\pi} \\ \frac{1}{4u^2}, & \textit{if } |u| > \frac{1}{2\pi}. \end{cases}$$

Proof: This is Lemma 2 in [3].

Lemma 2.3.

$$\begin{gathered} k'(0) = 0,\ k''(0) = \frac{\pi^8}{18}, \\ k'(\frac{1^+}{2\pi}) = -4\pi^3,\ k'(\frac{1^-}{2\pi}) = -4\pi^3 + \pi^5, \\ k''(\frac{1^+}{2\pi}) = 24\pi^4,\ k''(\frac{1^-}{2\pi}) = \frac{\pi^8}{2} - 4\pi^6 + 24\pi^4. \end{gathered}$$

Proof: Straightforward calculations.

Lemma 2.4.

$$\hat{k}(y) = -\frac{1}{(2\pi y)^2}\int_{-\infty}^{\infty} k''(u)e(-uy)du + \frac{\pi^3}{2y^2}\cos y,$$

where $\hat{f}$ is the Fourier transform of f,

$$\hat{f}(y) = \int_{-\infty}^{\infty} f(u)e(-uy)du,\ e(u) = e^{2\pi i u}.$$

Proof: Note that $k(u) \ll \min(1, 1/u^2)$, $k'(u) \ll \min(1, 1/u^3)$ and $k''(u) \ll \min(1, 1/u^4)$ except at $u = 1/2\pi$. Integrating by parts twice,

$$\begin{aligned}
\hat{k}(y) &= \int_{-\infty}^{\infty} k(u)e(-uy)du \\
&= \frac{-1}{2\pi i y}\int_{-\infty}^{\infty} k(u)de(-uy) = \frac{1}{2\pi i y}\int_{-\infty}^{\infty} e(-uy)k'(u)du \\
&= \frac{-1}{(2\pi i y)^2}\int_{-\infty}^{\infty} k'(u)de(-uy) = \frac{1}{(2\pi i y)^2}\int_{-\infty}^{\infty} e(-uy)dk'(u) \\
&= \frac{1}{(2\pi i y)^2}\int_{-\infty}^{\infty} k''(u)e(-uy)du \\
&\quad + \Big(k'(\frac{1^+}{2\pi}) - k'(\frac{1^-}{2\pi})\Big)e(\frac{y}{2\pi}) + \Big(k'(-\frac{1^+}{2\pi}) - k'(-\frac{1^-}{2\pi})\Big)e(-\frac{y}{2\pi}) \\
&= \frac{-1}{(2\pi y)^2}\int_{-\infty}^{\infty} k''(u)e(-uy)du + \frac{\pi^3}{2y^2}\cos y
\end{aligned}$$

by Lemma 2.3 and the fact that $k(u)$ is even.

Our key improvement is the following

Lemma 2.5. *Let $x = T^\beta$. For any $\beta > 0$,*

$$\begin{aligned}
&\sum_{0<\gamma,\gamma'\le T} \hat{k}((\gamma-\gamma')\log x)\frac{(\gamma-\gamma')^2}{4+(\gamma-\gamma')^2} \\
=&\frac{\pi^2 T}{16\log T}\frac{F(\beta)}{\beta^2} - \frac{T}{64\pi^4\log T\beta^3}\int_{-\infty}^{\infty} F(\alpha)k''\Big(\frac{\alpha}{2\pi\beta}\Big)d\alpha,
\end{aligned}$$

where $F(\alpha)$ is as defined in (2).

Proof: By Lemma 2.4, the above sum

$$\begin{aligned}
= &\quad \frac{\pi^3}{2(\log x)^2} \sum_{0<\gamma,\gamma'\le T} \cos\left((\gamma-\gamma')\log x\right)\frac{1}{4+(\gamma-\gamma')^2} \\
&- \frac{1}{(2\pi\log x)^2} \sum_{0<\gamma,\gamma'\le T} \int_{-\infty}^{\infty} k''(u)e(-u(\gamma-\gamma')\log x)du \frac{1}{4+(\gamma-\gamma')^2} \\
= &\quad \frac{\pi^3}{8(\log x)^2} \sum_{0<\gamma,\gamma'\le T} x^{i(\gamma-\gamma')} w(\gamma-\gamma') \\
&- \frac{1}{(4\pi\log x)^2} \int_{-\infty}^{\infty} \Big(\sum_{0<\gamma,\gamma'\le T} e(-u(\gamma-\gamma')\log x) w(\gamma-\gamma') \Big) k''(u)du \\
= &\quad \frac{\pi^3 \frac{T}{2\pi}\log T}{8(\log x)^2} F(\beta) - \frac{\frac{T}{2\pi}\log T}{(4\pi\log x)^2} \int_{-\infty}^{\infty} F(2\pi u\beta)k''(u)du
\end{aligned}$$

which gives the lemma by substituting $\alpha = 2\pi u\beta$ in the integral.

Using Lemma 2.2 and Lemma 2.5, we have the following improvement of Lemma 3 in [3]:

Lemma 2.6. *Let $x = T^\beta$. Then for $\beta > 0$,*

$$\begin{aligned}
R &= \frac{T}{(2\pi^2\beta)^2} \int_{-\infty}^{\infty} F(\alpha)k\Big(\frac{\alpha}{2\pi\beta}\Big)d\alpha + \frac{T}{16\log^2 T}\frac{F(\beta)}{\beta^3} \\
&\quad - \frac{T}{64\pi^6\beta^4\log^2 T} \int_{-\infty}^{\infty} F(\alpha)k''\Big(\frac{\alpha}{2\pi\beta}\Big)d\alpha + O(\log^3 T).
\end{aligned}$$

Proof: We simply note that

$$\begin{aligned}
\sum_{0<\gamma,\gamma'\le T} \hat{k}((\gamma-\gamma')\log x) &= \sum_{0<\gamma,\gamma'\le T} \hat{k}((\gamma-\gamma')\log x) w(\gamma-\gamma') \\
&\quad + \sum_{0<\gamma,\gamma'\le T} \hat{k}((\gamma-\gamma')\log x) \frac{(\gamma-\gamma')^2}{4+(\gamma-\gamma')^2}
\end{aligned}$$

Using Lemma 2.2 and Lemma 2.5, we get the second and third terms. Meanwhile,

$$\begin{aligned}
&\sum_{0<\gamma,\gamma'\le T} \hat{k}((\gamma-\gamma')\log x) w(\gamma-\gamma') \\
= &\int_{-\infty}^{\infty} k(u) \sum_{0<\gamma,\gamma'\le T} e(-u(\gamma-\gamma')\log x) w(\gamma-\gamma') du \\
= &\frac{T}{2\pi}\log T \int_{-\infty}^{\infty} F(2\pi u\beta)k(u)du \\
= &\frac{T\log T}{(2\pi)^2\beta} \int_{-\infty}^{\infty} F(\alpha)k\Big(\frac{\alpha}{2\pi\beta}\Big)d\alpha.
\end{aligned}$$

which accounts for the first term.

Lemma 2.7. *For any* $\beta > 0$,

$$\int_0^\beta T^{-2\alpha} k''\Big(\frac{\alpha}{2\pi\beta}\Big) d\alpha = 16\pi^2\beta^2(\log T)^2 \int_0^\beta T^{-2\alpha} k\Big(\frac{\alpha}{2\pi\beta}\Big) d\alpha.$$

Proof: Integrating by parts twice.

Lemma 2.8. *Assume the Riemann Hypothesis and Twin Prime Conjecture. For any* $\epsilon > 0$ *and* $0 < \beta < 1$,

$$\begin{aligned}\int_{-\infty}^{\infty} F(\alpha) k\Big(\frac{\alpha}{2\pi\beta}\Big) d\alpha \quad = \quad & 2\pi^2\beta^2\Big[1 - \frac{\pi^2}{8} + \log\frac{\pi}{2} + \int_1^\infty \frac{F(\alpha)}{\alpha^2} d\alpha - \log\beta\Big] \\ & + 2(\log T + C) \int_0^\beta T^{-2\alpha} k\Big(\frac{\alpha}{2\pi\beta}\Big) d\alpha \\ & + O\Big(\frac{1}{\beta^2 \log^4 T}\Big) + O\Big(\frac{\log T}{T^{(1/2-\epsilon)\beta}}\Big) + O\Big(\frac{\beta^2}{\log^2 T}\Big)\end{aligned}$$

with $C = -2\log 2\pi - 2$.

Proof: Let $\epsilon_T = 3\log\log T/\log T$. Since F and k are even,

$$\begin{aligned}\int_{-\infty}^{\infty} F(\alpha) k\Big(\frac{\alpha}{2\pi\beta}\Big) d\alpha =& 2\Big(\int_0^\beta + \int_\beta^{1-\epsilon_T} + \int_{1-\epsilon_T}^1 + \int_1^\infty\Big) F(\alpha) k\Big(\frac{\alpha}{2\pi\beta}\Big) d\alpha \\ =& 2(I_1 + I_2 + I_3 + I_4)\end{aligned}$$

From (3), with $C = -2\log 2\pi - 2$,

$$\begin{aligned} I_1 =& \int_0^\beta \Big[\alpha + \frac{logT + C}{T^{2\alpha}}\Big]\Big[\frac{\pi\beta}{\alpha} - \frac{\pi^2}{2}\cot\big(\frac{\pi\alpha}{2\beta}\big)\Big]^2 d\alpha \\ & + O\Big(\int_0^\beta \alpha T^{\alpha-1}\frac{\alpha^2}{\beta^2} d\alpha\Big) + O\Big(\int_0^\beta \frac{T^{-(1/2-\epsilon)\alpha}}{\log T}\frac{\alpha^2}{\beta^2} d\alpha\Big) \\ =& \int_0^\beta \alpha\Big[\frac{\pi\beta}{\alpha} - \frac{\pi^2}{2}\cot\big(\frac{\pi\alpha}{2\beta}\big)\Big]^2 d\alpha + (\log T + C)\int_0^\beta T^{-2\alpha} k\Big(\frac{\alpha}{2\pi\beta}\Big) d\alpha \\ & + O\Big(\frac{T^{\beta-1}}{\beta^2\log^4 T}\Big) + O\Big(\frac{1}{\beta^2\log^4 T}\Big)\end{aligned}$$

because $\cot(x) = 1/x + O(x)$ when $0 \le x \le \pi/2$. The first integral is elementary to evaluate.

$$I_1 = \pi^2\beta^2\Big[1 - \frac{\pi^2}{8} + \log\frac{\pi}{2}\Big] + (\log T + C)\int_0^\beta T^{-2\alpha} k\Big(\frac{\alpha}{2\pi\beta}\Big) d\alpha + O\Big(\frac{1}{\beta^2\log^4 T}\Big).$$

By (3) again, we have,

$$\begin{aligned}
I_2 =& \int_{\beta}^{1-\epsilon_T} \Big[\alpha + O(\alpha T^{\alpha-1}) + O\Big(\frac{\log T}{T^{(1/2-\epsilon)\alpha}}\Big)\Big]\Big(\frac{\pi\beta}{\alpha}\Big)^2 d\alpha \\
=&\pi^2\beta^2[\log(1-\epsilon_T) - \log\beta] + O\Big(\frac{1}{\log^4 T}\Big) + O\Big(\frac{\log T}{T^{(1/2-\epsilon)\beta}}\Big) \\
I_3 =& \int_{1-\epsilon_T}^{1} \Big[\alpha + O\Big(\frac{T^{\alpha-1}}{\log T}\Big)\Big]\Big(\frac{\pi\beta}{\alpha}\Big)^2 d\alpha \\
=& -\pi^2\beta^2\log(1-\epsilon_T) + O\Big(\frac{\beta^2}{\log T}\int_{1-\epsilon_T}^{1} T^{\alpha-1} d\alpha\Big) \\
=& -\pi^2\beta^2\log(1-\epsilon_T) + O\Big(\frac{\beta^2}{\log^2 T}\Big).
\end{aligned}$$

Finally,

$$I_4 = \pi^2\beta^2 \int_1^{\infty} \frac{F(\alpha)}{\alpha^2} d\alpha.$$

Combining the results for I_1, I_2, I_3 and I_4, we have the lemma.

Lemma 2.9. *Assume the Riemann Hypothesis and Twin Prime Conjecture. For any* $\epsilon > 0$ *and* $0 < \beta < 1$*, where* $\beta = \log x/\log T$*,*

$$\begin{aligned}
\int_{-\infty}^{\infty} F(\alpha) k''\Big(\frac{\alpha}{2\pi\beta}\Big) d\alpha =& 4\pi^6\beta^2 - 24\pi^4\beta^4 + 48\pi^4\beta^4 \int_1^{\infty} \frac{F(\alpha)}{\alpha^4} d\alpha \\
&+ 32\pi^2\beta^2(\log T)^2(\log T + C)\int_0^{\beta} T^{-2\alpha} k\Big(\frac{\alpha}{2\pi\beta}\Big) d\alpha \\
&+ O\Big(\frac{1}{\log^2 T}\Big) + O\Big(\frac{\log T}{T^{(1/2-\epsilon)\beta}}\Big)
\end{aligned}$$

Proof: Let $\epsilon_T = 3\log\log T/\log T$. Again, since F and k are even,

$$\begin{aligned}
\int_{-\infty}^{\infty} F(\alpha) k''\Big(\frac{\alpha}{2\pi\beta}\Big) d\alpha =& 2\Big(\int_0^{\beta} + \int_{\beta}^{1-\epsilon_T} + \int_{1-\epsilon_T}^{1} + \int_1^{\infty}\Big) F(\alpha) k''\Big(\frac{\alpha}{2\pi\beta}\Big) d\alpha \\
=& 2(J_1 + J_2 + J_3 + J_4)
\end{aligned}$$

By (3),

$$\begin{aligned}
J_1 =& \int_0^{\beta} \alpha k''\Big(\frac{\alpha}{2\pi\beta}\Big) d\alpha + 16\pi^2\beta^2(\log T)^2(\log T + C)\int_0^{\beta} T^{-2\alpha} k\Big(\frac{\alpha}{2\pi\beta}\Big) d\alpha \\
&+ O\Big(\int_0^{\beta} \alpha T^{\alpha-1} d\alpha\Big) + O\Big(\int_0^{\beta} \frac{T^{-(1/2-\epsilon)\alpha}}{\log T} d\alpha\Big) \\
=& 4\pi^2\beta^2 \int_0^{1/2\pi} u k''(u) du + 16\pi^2\beta^2(\log T)^2(\log T + C)\int_0^{\beta} T^{-2\alpha} k\Big(\frac{\alpha}{2\pi\beta}\Big) d\alpha \\
&+ O\Big(\frac{T^{\beta-1}}{\log^2 T}\Big) + O\Big(\frac{1}{\log^2 T}\Big)
\end{aligned}$$

since $k''(x) \ll 1$ when $0 \le x \le 1/2\pi$. Using integration by parts twice and Lemma 2.3 to compute the first integral, we have

$$\begin{aligned} J_1 =& 2\pi^4(\pi^2-6)\beta^2 + 16\pi^2\beta^2(\log T)^2(\log T + C)\int_0^\beta T^{-2\alpha} k\Big(\frac{\alpha}{2\pi\beta}\Big) d\alpha \\ &+ O\Big(\frac{1}{\log^2 T}\Big). \end{aligned}$$

By (3) again, we have

$$\begin{aligned} J_2 =& \int_\beta^{1-\epsilon_T} \Big[\alpha + O(\alpha T^{\alpha-1}) + O\Big(\frac{\log T}{T^{(1/2-\epsilon)\alpha}}\Big)\Big]\Big(\frac{24\pi^4\beta^4}{\alpha^4}\Big) d\alpha \\ =& 12\pi^4\beta^4\Big[\frac{1}{\beta^2} - \frac{1}{(1-\epsilon_T)^2}\Big] + O\Big(\frac{1}{\log^4 T}\Big) + O\Big(\frac{\log T}{T^{(1/2-\epsilon)\beta}}\Big) \\ J_3 =& \int_{1-\epsilon_T}^1 \Big[\alpha + O\Big(\frac{T^{\alpha-1}}{\log T}\Big)\Big]\Big(\frac{24\pi^4\beta^4}{\alpha^4}\Big) d\alpha \\ =& 12\pi^4\beta^4\Big[\frac{1}{(1-\epsilon_T)^2} - 1\Big] + O\Big(\frac{\beta^4}{\log T}\int_{1-\epsilon_T}^1 T^{\alpha-1} d\alpha\Big) \\ =& 12\pi^4\beta^4\Big[\frac{1}{(1-\epsilon_T)^2} - 1\Big] + O\Big(\frac{\beta^4}{\log^2 T}\Big). \end{aligned}$$

Finally,

$$J_4 = 24\pi^4\beta^4 \int_1^\infty \frac{F(\alpha)}{\alpha^4} d\alpha.$$

Combining the results for J_1, J_2, J_3 and J_4, we have the lemma.

Combining Lemma 2.6, Lemma 2.8 and Lemma 2.9, we have

Lemma 2.10. *Assume the Riemann Hypothesis and Twin Prime Conjecture, for fixed* $0 < \beta < 1$*, where* $\beta = \log x/\log T$*,*

$$\begin{aligned} R =& \frac{T}{2\pi^2}\Big[1 - \frac{\pi^2}{8} + \log\frac{\pi}{2} + \int_0^\infty \frac{F(\alpha)}{\alpha^2} d\alpha - \log\beta\Big] + \frac{3T}{8\pi^2\log^2 T} \\ &- \frac{3T}{4\pi^2\log^2 T}\int_1^\infty \frac{F(\alpha)}{\alpha^4} d\alpha + O\Big(\frac{T}{\log^2 T}\Big) + O\Big(\frac{T}{\beta^4\log^4 T}\Big) \end{aligned}$$

Note: This is a more precise version of Lemma 4 of [3]. Also, we keep some of the $T/\log^2 T$ terms explicit because one can actually make the $O(T/\log^2 T)$ error term $= C_1 T/\log^2 T + O(T\log\log T/\log^3 T)$ for some constant C_1 by using Theorem 1.1 in [2].

Following [3], we need to compute the mean value of the Dirichlet series in Lemma 2.1, and the cross term obtained from multiplying $S(t)$ with this series. Let

$$G(T) = \int_1^T \Big|\frac{1}{\pi}\sum_{n\le x} \frac{\Lambda(n)}{n^{1/2}} \frac{\sin(t\log n)}{\log n} f\Big(\frac{\log n}{\log x}\Big)\Big|^2 dt$$

and

$$H(T) = \frac{2}{\pi} \int_1^T S(t) \sum_{n \le x} \frac{\Lambda(n)}{n^{1/2}} \frac{\sin(t \log n)}{\log n} f\Big(\frac{\log n}{\log x}\Big) dt,$$

where f is defined as in (5). We need a lemma.

Lemma 2.11. *For $C \ge 2$ and $k \ge 1$,*

$$\sum_{n=1}^{\infty} \frac{n^k}{C^n} \ll_k \frac{1}{C}.$$

Proof: First, we note that $u^k C^{-u}$ is decreasing when $u > \frac{k}{\log C}$. So,

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{n^k}{C^n} &= \sum_{n=1}^{k/\log C} \frac{n^k}{C^n} + \sum_{n > k/\log C} \frac{n^k}{C^n} \\ &\le \Big(\frac{k}{\log 2}\Big)^k \frac{1}{C-1} + \int_1^{\infty} u^k C^{-u} du \\ &\ll_k \frac{1}{C} + \frac{1}{\log C} C^{-1} \end{aligned}$$

by integration by parts. This gives the lemma.

From p.165-166 of [3], we have, assuming the Riemann Hypothesis,

$$G(T) = \frac{T}{2\pi^2} \sum_{n \le x} \frac{\Lambda^2(n)}{n \log^2 n} f^2\Big(\frac{\log n}{\log x}\Big) + O(x^2), \tag{6}$$

$$H(T) = -\frac{T}{\pi^2} \sum_{n \le x} \frac{\Lambda^2(n)}{n \log^2 n} f\Big(\frac{\log n}{\log x}\Big) + O(x^{2+\epsilon}) \tag{7}$$

for any $\epsilon > 0$. Adding (6) and (7), we have

$$\begin{aligned} G(T) + H(T) =& \frac{T}{2\pi^2} \Big[\sum_{p \le x} \frac{1}{p} f^2\Big(\frac{\log p}{\log x}\Big) - 2 \sum_{p \le x} \frac{1}{p} f\Big(\frac{\log p}{\log x}\Big) - \sum_{m=2}^{\infty} \sum_{p^m \le x} \frac{1}{m^2 p^m} \\ &+ \sum_{m=2}^{\infty} \sum_{p^m \le x} \frac{1}{m^2 p^m} \Big(f\Big(\frac{m \log p}{\log x}\Big) - 1\Big)^2 \Big] + O(x^{2+\epsilon}) \\ =& \frac{T}{2\pi^2} [S_1 - 2S_2 - S_3 + S_4] + O(x^{2+\epsilon}). \end{aligned}$$

$$\begin{aligned} S_3 &= \sum_{m=2}^{\infty} \sum_p \frac{1}{m^2 p^m} + O\Big(\sum_{m=2}^{\infty} \frac{1}{m^2} \sum_{n \ge x^{1/m}} \frac{1}{n^m} \Big) \\ &= \sum_{m=2}^{\infty} \sum_p \frac{1}{m^2 p^m} + O\Big(\sum_{m=2}^{\infty} \frac{1}{m^2 (m-1) x^{1-1/m}} \Big) \\ &= \sum_{m=2}^{\infty} \sum_p \frac{1}{m^2 p^m} + O\Big(\frac{1}{x^{1/2}} \Big). \end{aligned}$$

By Taylor's expansion of $\tan x$, we have $f(u) = 1 + O(u^2)$ when $0 \le u \le 1$. Thus,

$$\begin{aligned} S_4 &\ll \frac{1}{\log^4 x} \sum_{m=2}^{\infty} m^2 \sum_{p^m \le x} \frac{\log^4 p}{p^m} \\ &\ll \frac{1}{\log^4 x} \sum_{m=2}^{\infty} m^2 \sum_{i=1}^{\infty} \sum_{2^i \le p \le 2^{i+1}} \frac{i^4}{2^{mi}} \\ &\ll \frac{1}{\log^4 x} \sum_{m=2}^{\infty} m^2 \sum_{i=1}^{\infty} \frac{i^4}{(2^{m-1})^i} \\ &\ll \frac{1}{\log^4 x} \sum_{m=2}^{\infty} \frac{m^2}{2^{m-1}} \ll \frac{1}{\log^4 x} \end{aligned}$$

by using Lemma 2.11 twice.

We now define

$$T(u) = \sum_{2 \le p \le u} \frac{1}{p},$$

and have

$$T(u) = \log\log u + C_0 + \sum_p \Big(\log\Big(1 - \frac{1}{p}\Big) + \frac{1}{p}\Big) + r(u),$$

where $r(u) \ll \log u/\sqrt{u}$ under the Riemann Hypothesis, and the sum over primes on the right is equal to

$$-\sum_{m=2}^{\infty} \sum_p \frac{1}{mp^m}.$$

Then,

$$\begin{aligned} S_1 &= \int_2^x f^2\Big(\frac{\log u}{\log x}\Big) dT(u) \\ &= \int_2^x f^2\Big(\frac{\log u}{\log x}\Big) \frac{du}{u \log u} + \int_2^x f^2\Big(\frac{\log u}{\log x}\Big) dr(u) = I_1 + I_2. \end{aligned}$$

Similarly,

$$S_2 = \int_2^x f\Big(\frac{\log u}{\log x}\Big) \frac{du}{u \log u} + \int_2^x f\Big(\frac{\log u}{\log x}\Big) dr(u) = J_1 + J_2.$$

Thus,

$$S_1 - 2S_2 = I_1 - 2J_1 + (I_2 - 2J_2).$$

By integration by parts,

$$\begin{aligned}
I_2 - 2J_2 &= -r(2^-)\Big[f^2\Big(\frac{\log 2}{\log x}\Big) - 2f\Big(\frac{\log 2}{\log x}\Big)\Big] - \int_2^x r(u) \\
&\quad \Big[2f\Big(\frac{\log u}{\log x}\Big)f'\Big(\frac{\log u}{\log x}\Big) - f'\Big(\frac{\log u}{\log x}\Big)\Big]\frac{du}{u\log x} \\
&= r(2^-) - r(2^-)\Big[f\Big(\frac{\log 2}{\log x}\Big) - 1\Big]^2 \\
&\quad -\frac{2}{\log x}\int_2^x r(u)f'\Big(\frac{\log u}{\log x}\Big)\Big[f\Big(\frac{\log u}{\log x}\Big) - 1\Big]\frac{du}{u} \\
&= -\log\log 2 - C_0 + \sum_{m=2}^{\infty}\sum_p \frac{1}{mp^m} + O\Big(\frac{1}{\log^4 x}\Big) \\
&\quad +O\Big(\frac{1}{\log x}\int_2^x \frac{\log u}{\sqrt{u}}\Big(\frac{\log u}{\log x}\Big)\Big(\frac{\log u}{\log x}\Big)^2\frac{du}{u}\Big) \\
&= -\log\log 2 - C_0 + \sum_{m=2}^{\infty}\sum_p \frac{1}{mp^m} + O\Big(\frac{1}{\log^4 x}\Big)
\end{aligned}$$

because $f(u) = 1 + O(u^2)$ and $f'(u) \ll u$ when $0 \le u \le 1$. The integrals in I_1 and J_1 are elementary to evaluate. Using $u\cot u = 1 - u^2/3 + O(u^4)$ and $\sin u = u - u^3/6 + O(u^5)$, one has

$$\begin{aligned}
I_1 &= \log\log x - \log\log 2 - \frac{\pi^2}{8} + 1 - \log\frac{\pi}{2} + \frac{\pi^2(\log 2)^2}{12\log^2 x} + O\Big(\frac{1}{\log^4 x}\Big), \\
J_1 &= \log\log x - \log\log 2 - \log\frac{\pi}{2} + \frac{\pi^2(\log 2)^2}{24\log^2 x} + O\Big(\frac{1}{\log^4 x}\Big).
\end{aligned}$$

Hence,

$$S_1 - 2S_2 = -\log\log x + \log\frac{\pi}{2} - \frac{\pi^2}{8} + 1 - C_0 + \sum_{m=2}^{\infty}\sum_p \frac{1}{mp^m} + O\Big(\frac{1}{\log^4 x}\Big).$$

Therefore,

$$\begin{aligned}
G(T) + H(T) =&\frac{T}{2\pi^2}\Big[-\log\log x + \log\frac{\pi}{2} - \frac{\pi^2}{8} + 1 - C_0 \\
&+ \sum_{m=2}^{\infty}\sum_p\Big(\frac{1}{m} - \frac{1}{m^2}\Big)\frac{1}{p^m}\Big] + O\Big(\frac{T}{\log^4 x}\Big).
\end{aligned} \tag{8}$$

3 Proof of Theorem 1.1

Suppose $x = T^\beta$ and β is a fixed positive number less than $1/2$. We have, by Lemma 2.1, that (4) holds except on a countable set of points. Hence, on squaring both sides of (4) and integrating from 1 to T,

$$\int_1^T (S(t))^2 dt + H(T) + G(T) = R + O(T^{1/2} x^{1/2}),$$

where the error term is obtained by Cauchy-Schwarz inequality since $R \ll T$. The lower limit of integration may be replaced by zero since $\int_0^1 (S(t))^2 dt \ll 1$. Then, Lemma 2.10 and (8) give the theorem. The author would like to thank Professor Daniel Goldston for suggestion and discussion on this problem.

References

[1] T.H. Chan, *On a conjecture of Liu and Ye*, submitted.

[2] T.H. Chan, *More precise Pair Correlation Conjecture on the zeros of the Riemann zeta function*, submitted.

[3] D.A. Goldston, *On the Function $S(T)$ in the Theory of the Riemann Zeta-Function*, J. Number Theory **27** (1987), 149-177.

[4] H.L. Montgomery, *The pair correlation of zeros of the zeta function*, Analytic Number Theory (St. Louis Univ., 1972), Proc. Sympos. Pure Math. **24**, Amer. Math. Soc., Providence, 1973, pp. 181-193.

Tsz Ho Chan
Case Western Reserve University
Mathematics Department, Yost Hall 220
10900 Euclid Avenue
Cleveland, OH 44106-7058
USA
txc50@po.cwru.edu